

Synthesis of Cyclic Polybutadiene via Ring-Opening Metathesis Polymerization: The Importance of Removing Trace Linear Contaminants

Christopher W. Bielawski, Diego Benitez, and Robert H. Grubbs

Arnold and Mabel Beckman of Laboratory Synthesis, Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA 91125.

Supporting Information

Table 1. Synthesis of polybutadienes using 1,5-cyclooctadiene as monomer^a

[M/C] ₀	[Monomer] ₀ (M)	% Yield ^b	M _n ^c (x10 ³ Da)	PDI ^c
25	1.0	84	2.3	1.6
50	1.0	86	5.0	1.7
100	1.0	88	8.5	1.7
1150	1.0	87	86	1.9
100	0.5	83	2.2	1.7
100	0.1	0 ^e	ND	ND
100	4.0	90	26	1.9
100	8.1 ^f	95	145	1.8

^a Polymerizations were conducted in CH₂Cl₂ at 45 °C for 12 h. Monomer (M) = 1,5-cyclooctadiene (COD). Catalyst (C) = **1**. ^b Isolated yields. ^c Determined by size exclusion chromatography (SEC) with CH₂Cl₂ as eluent. The values are reported in Da and are relative to monodispersed polybutadiene standards. ^e Only low MW oligomers were observed by SEC. ^f Polymerization was performed in bulk COD. ND = Not determined.

Table 2. Synthesis of polybutadienes using 1,5,9-*trans-cis-trans*-cyclododecatriene as monomer^a

[M/C] ₀	[Monomer] ₀ (M)	% Yield ^b	M _n ^c (x10 ³ Da)	PDI ^c
1000	2.5	85	12	1.9
2500	2.5	80	32	1.8
6000	5.0	90	88	1.7
2500	1.0	0 ^e	ND	ND

^a Polymerizations were conducted in CH₂Cl₂ at 45 °C for 12 h. Monomer (M) = 1,5,9-*trans-cis-trans*-cyclododecatriene (CDT). Catalyst (C) = **1**. ^b Isolated yields. ^c Determined by size exclusion chromatography (SEC) with CH₂Cl₂ as eluent. The values are reported relative to monodispersed polybutadiene standards. ^e Only low MW oligomers were observed by SEC. ND = Not determined.